

# Neurologic Abnormalities in the Skeletal Dysplasias: A Clinical and Radiological Perspective

Ralph S. Lachman\*

International Skeletal Dysplasia Registry, Cedars-Sinai Medical Center, University of California Los Angeles and Harbor/UCLA Medical Center, Los Angeles

The neurologic manifestations of the skeletal dysplasias are reviewed. Three important major groups are identified: *Achondroplasia* (cranio-cervical junction problems in infancy, spinal stenosis and neurogenic claudication in the adult), *Type II collagenopathies* (upper cervical spine anatomic and functional problems), and *craniotubular and sclerosing bone dysplasias* (osseous overgrowth with foraminal obstruction problems). The remainder of the well-identified 150 or so bone dysplasias are also evaluated in depth for their diverse neurologic abnormalities. The findings discussed are important both for the diagnosis and management of these patients. *Am. J. Med. Genet.* 69:33–43, 1997. © 1997 Wiley-Liss, Inc.

**KEY WORDS:** skeletal dysplasias; bone dysplasias; osteochondrodysplasias; neurologic abnormalities

## INTRODUCTION

The most important neurological abnormalities in the skeletal dysplasias are the central nervous system (CNS) and high cervico-medullary problems in *achondroplasia* and the upper cervical spine complications in the *Type II collagenopathies* and their related disorders. However, a large variety of neurological associations and manifestations may occur in many of the 150 or so well-defined skeletal dysplasias. They are impor-

tant not only for diagnostic purposes, but also at times for clinical management.

## ACHONDROPLASIA

Achondroplasia is due to a defect in enchondral bone formation with resulting decrease in growth, not only of long bones but also a decrease in size of the vertebral bodies and of the diameter of the posterior vertebral arches. The foramen magnum is also affected with resultant decrease in its size and alterations in its shape, especially at the expense of its lateral walls. Secondly, there is a potential for over growth of membranous bone (the greatest portion of the skull convexity).

The early life (infancy) neurologic complications of achondroplasia are found at the cranio-cervical junction and relate to the "small foramen magnum"—an oxymoron. Secondary *significant hydrocephalus* may occur but is infrequent, it is communicating, associated with intracranial venous hypertension, megalencephaly and medullary, and upper cervical cord compression. The cranio-cervical compression can exist independent of the hydrocephalus and may manifest as sleep apnea and "aborted" or "real" Sudden Infant Death "syndrome," apneustic breathing [Mador et al., 1990], and increasing upper limb weakness. A recent study suggests that the best predictors of the need for suboccipital decompression include the demonstration of lower limb hyperreflexia (or clonus); central hypopnea (apnea) at poly-somnography; and reduced foramen magnum measurements [Pauli et al., 1995].

Radiographic evaluation should include: (1) CT of the foramen magnum (size) and brain (hydrocephalus), and (2) MRI of the brain and upper cervical-medullary region, including CSF flow studies across the foramen. MRI performed in the flexed neck position may also contribute important information.

Other abnormalities detected on imaging include: upper cervical cord thinning secondary to compression and cord syrinx following infarction/hemorrhage (Figs. 1–5).

Recently, a nontraumatic cervical cord infarction in a 12-year-old with resultant quadriplegia has been reported [Wieting and Krach, 1994]. It is apparent that the achondroplastic patient is "not out of the woods" for this complication even in later life.

Usually later in life (young/late adult), nerve root compression (spinal stenosis) occurs as well as inter-

Contract grant sponsor: NIH; Contract grant number: PO1 HD22657-08.

The author submits this manuscript in honor of the 65th birthday of Jürgen Spranger, colleague, pioneer, discerning spirit, but most of all—a good friend. I wish him many more wonderful years with us among the skeletal dysplasias.

Wenn man einen Zwerg gesehen hat, hat man alle gesehen (if you have seen one dwarf, you've seen them all). He as much as anyone has shown us the absolute *untruth* in that statement.

\*Correspondence to: Ralph S. Lachman, M.D., International Skeletal Dysplasia Registry, Cedars-Sinai Medical Center, Room 1001, 444 S. San Vicente Blvd., Los Angeles, CA 90048.

Received 10 January 1996; Accepted 11 June 1996




Fig. 1. Six-month-old achondroplasia patient. **A, B:** CT of the skull revealing decreased side wall diameter of the foramen magnum and oval shape to the foramen magnum. In contrast, there is a normal C<sub>1</sub>C<sub>2</sub> diameter and shape at the odontoid level.

mittent (or persistent) neuron claudication, most commonly in the lumbar spine resulting in parasthesias and progressive paraparesis and even rarely quadriplegia [Fortuna et al., 1989].

Imaging of these problems includes: (1) plain films (AP and lateral) of the spine, and (2) MRI of spine area in question. Imaging findings may include: (1) spinal stenosis, (2) hypoplastic T12/L1 vertebral body, (3) herniated discs (multiple sites), and (4) nerve entrapment.

The *clinical situation* must be evaluated as well as the imaging studies to decide if laminectomy should be performed. Postlaminectomy recurrences are not rare. Two much rarer neurologic entities described in achondroplasia are tethered cord [Phadke et al., 1990], and brain tumor [1 case] [McArdle et al., 1984].

### TYPE II COLLAGENOPATHIES

Most of the type II collagenopathies manifest abnormalities in the cervical spine (Table I). These abnor-




Fig. 2. Same patient as Figure 1. **A, B:** MRI of the brain stem and cervical spine with measurements of both anterior-posterior and lateral dimensions of the foramen magnum. There is no evidence of cervical cord compression.




Fig. 3. Same patient as Figures 1 and 2. **A, B, C:** CT of the brain showing mild/moderate ventricular dilatation.

malities include C1-C2 subluxation/dislocation secondary to abnormal formation of the odontoid process, the base of C2, ossification of C1, and abnormally lax or non-fixed posterior portion of the ligament extending from C1 to C2. Any or all of the abnormalities can result in the abnormal posterior displacement of C2 into the neural canal, especially with extreme flexion. Clinically, this problem can manifest throughout life, but even in infancy it may occur especially if the patient needs intubation (i.e., for anesthesia) or suffers trauma and the neck is acutely flexed. All patients with a type II collagenopathy, except for Stickler dysplasia, should be screened for this problem and patients with unclassifiable forms of SED [Anderson et al., 1982; LeDoux et al., 1991; Merrill et al., 1989; Yagi et al., 1987].

Imaging studies should include: (1) AP and lateral cervical spine (2) lateral flexion/extension views of the cervical spine, (3) CT (with/without) reconstruction of the C-spine for anatomic delineation, if necessary, and (4) MRI of the cervical cord region for possible cord damage. One should note that the most important studies for evaluation are 2 and 4 (Figs. 6–8).

Recently, doubt has been raised about the frequency of this problem in patients with some forms of SED (SEMD), which has not been the experience of most [Reardon et al., 1994]. Our experience suggests that those with a pectus carinatum (high sternum) are protected by the inability to flex significantly.

As one goes through the remainder of the 150 well-defined skeletal dysplasias, it is apparent that there is




Fig. 4. Four-month-old infant with achondroplasia. **A, B:** MRI of the cervical cord and brainstem (pre-operatively) with a syrinx of the cervical cord and cervical cord compression at the medullary-cord junction.

a large variety of other neurologic abnormalities, associations, and complications. Many of these are related to the skeletal changes themselves. The less important ones are covered in Table II.

#### CAMPOMELIC DYSPLASIA

This entity may or may not be lethal and commonly has mild or severe CNS abnormalities, which can present with generalized hypotonia. Severe cervical kyphosis may lead to cervical cord problems [Coscia et al., 1989].

#### Chondrodysplasia Punctata (CP) Group of Disorders

**CP—Rhizomelic type.** This peroxisomal enzyme abnormality disorder often results in perinatal or early infancy death. Survivors may manifest: facial paralysis, spastic quadriplegia, generalized spasticity, and severe psychomotor retardation [Lenti et al., 1991; Williams et al., 1991]. They appear to be in chronic pain [Rimoin, 1996].

**CP—Conradi-Hunermann type.** Four cases of progressive myelopathy have been described [Goodman et al., 1990].

**CP—Brachytelephalangic type.** Hypotonia has been a prominent symptom and sign (Fig. 9).

#### Cranio-Tubular Dysplasias and Sclerosing Bone Dysplasias

In several of these dysplasias, there is osseous overgrowth of the skull base and encroachment of the foramina resulting in neurologic and neurovascular problems [Kirkpatrick et al., 1977].

**Craniodiaphyseal dysplasia.** Increasing deafness, vision loss and sudden death, probably on a vascular basis.

**Craniometaphyseal dysplasia.** Hydrocephalus, optic atrophy, progressive deafness, facial nerve palsy, hemiplegia, quadraplegia, and medullary compression have been noted [Allen et al., 1982; Hudgins et al., 1987; Saper et al., 1974].

**Diaphyseal dysplasia, Engelmann type.** Increased intracranial pressure (hydrocephalus), progressive hearing loss, and visual loss may occur [Applegate et al., 1991; Paludetti et al., 1994].

**Dysosteosclerosis.** Beginning in adolescence, decreased visual acuity, optic atrophy, and facial palsy [Chitayat et al., 1992]. These findings are not seen in the dominant form of osteopetrosis.

**Endosteal hyperostosis, Van Buchem.** Hydrocephalus (increased intracranial pressure) and rarely spinal cord compression may occur. Cranial nerve palsies, especially of the facial nerves, are not infrequent [Fryns et al., 1988; Kitasawa et al., 1986].






Fig. 5. Same patient as in Figure 4, 2 months later. The postoperative MRI study shows the surgically induced posterior ossification defects producing increased space for the cord.

**Endosteal hyperostosis, Worth.** Mild facial, optic, and auditory nerve involvement may occur with progression in the adult (5 cases). Chronic increased intracranial pressure and cerebellar tonsillar herniation into the foramen magnum were reported recently [Adès et al., 1994].

**Frontometaphyseal dysplasia.** Progressive hearing loss has occurred in a few patients [Arenberg et al., 1974].

**Osteopathia striata with cranial sclerosis.** Facial nerve palsy can be seen as well as other mild cranial nerve abnormalities such as deafness and ophthalmoplegia [Gay et al., 1994].

TABLE I. Type II Collagenopathies

Name	Cervical spine problems
Achondrogenesis II	Lethal
Hypochondrogenesis	Yes
Spondylo-epiphyseal dysplasia (SED) congenita	Yes
Spondylo-epi-metaphyseal dysplasia (SEMD) Strudwick	Yes
Kniest dysplasia	Yes
Stickler dysplasia <sup>a</sup>	No
Autosomal dominant spondyloarthropathy (mild SED)	?
Goldblatt syndrome (SEMD with dentinogenesis imperfecta and joint laxity)	?
SED-Namaqualand	?
SED-Capetown	?

<sup>a</sup> Cervical myelopathy secondary to spinal stenosis (Noel).

**Osteopetrosis.** The severe autosomal recessive form can manifest with multiple cranial nerve palsies, including facial, optic and auditory involvement in infancy [Abdel-al et al., 1994; Al-Mefty et al., 1988; Bartgaski et al., 1989; Benecke et al., 1993; Demirci et al., 1991; Elster et al., 1992; Wilms et al., 1990]. Myelopathy secondary to cervical spine involvement can occur [McCleary et al., 1987].

The autosomal dominant form has presented with carpal tunnel syndrome [Rakic et al., 1986].

**Sclerosteosis.** Cranial nerve palsies especially involving the seventh nerve can be seen as well as




Fig. 6. Three-year-old with spondyloepiphyseal dysplasia (SED) congenita. The MRI of the cervical region reveals C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub> posterior subluxation with cord compression in the neutral position.




Fig. 7. Seven-year-old with SED congenita. **A, B:** MRI shows typical C<sub>1</sub>C<sub>2</sub> subluxation with cord impingement.

increased intracranial pressure (hydrocephalus) [Dort et al., 1990; duPlessis et al; 1993; Hill et al., 1986].

#### DIASTROPHIC DYSPLASIA

There is significant cervical kyphosis in about 1/3 of patients, with resultant cervical cord compression in some. Quadriplegia may occur especially related to

anesthesia [Krecak et al., 1987; Lachman et al., 1981; Richards et al., 1991].

#### DYGGVE-MELCHIOR-CLAUSSEN DYSPLASIA

The more severely involved cases may exhibit progressive basilar impression and even brain death [Spranger et al., 1976].




Fig. 8. Same patient as in Figure 7. CT reconstruction reveals the odontoid hypoplasia with an Os odontoideum. The measurement of the anterior-posterior cervical spine dimension and posterior dislocation of C<sub>2</sub> is apparent.

TABLE II. Other Skeletal Dysplasias With Neurologic Abnormalities

Skeletal dysplasia	Abnormality	Frequency	References
Acrodysostosis	Psychomotor retardation Lower extremity neuropathy (lumbar stenosis, tight dural sac) Optic atrophy, strabismus, seizures and choreoathetosis Tapetoretinal degeneration cerebellar ataxia Syrinx of the cervical cord Hydrocephalus Agenesis of the corpus callosum Dandy Walker malformation Forebrain abnormalities with encephalocele	77% of cases Uncommon  Rare Major manifestation Uncommon 2 cases Rare Uncommon  Frequent	Taybi, Lachman, 1996 Hamanshi, 1993  Ellis, 1984 Popovic-Rolovic, 1976 personal experience Trabelsii, 1990  Chervenak, 1986
Acrodysplasia with retinitis pigmentosa and nephropathy (Saldino-Mainzer dysplasia) Acromesomelic dysplasia			
Asphyxiating thoracic dysplasia			
Atelosteogenesis type I			
Boomerang dysplasia Cleidocranial dysplasia Desbuquois dysplasia	Encephalocele Syringomyelia Sleep apnea and sudden infant death syndrome (SIDS) with cervical kyphosis	? 3 cases Several cases	Canki-Klain, 1992 Dore, 1987 Sillence, 1995 Beemer, 1994
Dyssegmental dysplasia-Rolland-Desbuquois type Silverman-Handmaker type Lowry Wood dysplasia (MED, Lowry-Wood) Metaphyseal chondrodysplasia, Swachman-Diamond type	Hydrocephalus Dandy Walker cyst Hypoplasia of corpus callosum Focal pontine leuodystrophy	Common Common 1 case 3 cases	Aleck, 1987  Yamamoto, 1995 Andres, 1993 Mah, 1987 Steinsapir, 1985 Rosenberg, 1986
Metaphyseal-Sella turcica dysplasia (Rosenberg)	Thickening of sella turcica without clinical symptomatology	?	
Multiple exostosis	Pain, parasthesias, and motor problems secondary to exophytic impingement	Common	N'dri, 1994 O'Brien, 1994 Zonana, 1977
Opsismodysplasia (Fig. 10)	Hyptonia, C1C2 dislocation and cord compression with severe ossification defect	Common	
Osebold Remondini dysplasia Osteopoikilosis Pseudodiastrophic dysplasia Pycnodysostosis	Numbness and dythesias (pathogenesis unclear) Lumbar spine stenosis with clinical symptomatology Hydrocephalus Sleep apnea (medullary compression-bony overgrowth) Hangman fractures (spondylolysis of cervical spine) Porencephalic cyst Severe brain abnormalities (arrhinencephaly, vermis hypoplasia, arachnoid cysts and cerebral dysgenesis) Optic atrophy (chiasm and nerves) C <sub>1</sub> C <sub>2</sub> subluxation	? ? 1 case ? Uncommon 1 case Common  ? 1 case	Taybi, Lachman, 1996 Weisz, 1982 Taybi, Lachman, 1996 Floman, 1989 Currarino, 1989 Figueiredo, 1989 Pradio, 1993  Cideciyan, 1993 Taybi, Lachman, 1996
Short rib polydactyly dysplasia Type II and IV Type IV Spondylometaphyseal dysplasias (SMD) SMD-corner fracture (Schmidt-Sutcliffe) type SMD-Sedeghat type SMD-Frydman type Tricho-rhino phalangeal dysplasia, type I	Lissencephaly Quadripareisis and basal ganglia calcifications Mental retardation and seizure disorder	? ? Very common	
Weissenbacher-Zweymuller dysplasia Walcott Rallison dysplasia (SED with Diabetes Mellitus) Yunis Varon dysplasia	Neural tube defect Seizures, unrelated to hypoglycemia Agenesis of the corpus callosum	2 cases (Monozygotic twins) 1 case 1 case	Hamers, 1985 Balza, 1990 Ramer, 1993 Taybi, Lachman, 1996  Taybi, Lachman, 1996




Fig. 9. Four-year-old with an unclassifiable form of chondrodysplasia punctata. An MRI of the cervical region shows cervical spine (C<sub>3</sub>) subluxation with cord compression.

### MAFFUCCI (SYNDROME) DYSPLASIA

This entity of multiple enchondromatosis and vascular abnormalities may have cranial nerve involvement when the base of the skull is affected. There have been five cases of brain tumors (gliomas) reported, a very high incidence for this rare entity [Nakayama et al., 1994; Robinson et al., 1994].

### METATROPIC DYSPLASIA

Severe upper cervical spine problems are often present, especially C<sub>1</sub> C<sub>2</sub> subluxations/dislocations secondary to either hypoplastic or enlarged odontoid and/or a malformed base of C<sub>2</sub> resulting in sleep apnea and other symptoms similar to the type II collagenopathies. These patients definitely should be screened [Shohat et al., 1989]. Hydrocephalus also has been identified, as well as a recent report of a primary motor and sensory neuropathy with storage abnormalities in the rough endoplasmic reticulum of the nerve fibers similar to what is seen in the growth plate of bones [Yamashita et al., 1994].

### OSTEOGENESIS IMPERFECTA (OI)

In the nonlethal OI type III, hydrocephalus with macrocrania is sometimes present [Charnas et al.,



Fig. 10. Five-year-old with opsismodysplasia. On plain films of the cervical spine, one sees a severe lack of ossification of the cervical vertebral bodies in this rare disorder.

1993; Pauli et al., 1986; Rush et al., 1984; Tsipouras et al., 1986]. Recently, clinical manifestations secondary to basilar impression have been identified primarily in OI type IV [Sillence et al., 1994]. Rupture of a cerebral aneurysm in a fenestrated vertebral artery perhaps secondary to the underlying type I collagen defect also has been reported in OI type I (1 case). Reflex sympathetic dystrophy can occur [Karras et al., 1993].

### THANATOPHORIC DYSPLASIA

This is probably the most common lethal skeletal dysplasia, with an incidence of almost 2 per 100,000 pregnancies. All cases with neuropathologic examination show severe brain abnormalities, including abnormal sulci with polymicrogyria and neuronal heterotopia and also marked hypoplasia of the middle and posterior cranial fossae. This suggests severe arrest of cerebral cortical development [Coulter et al., 1991]. Early extreme hydrocephalus with craniosynostosis is the pathogenesis in the group with Kleeblattschädel (cloverleaf skull). Rare survivors with thanatophoric dysplasia are all very severely neurologically impaired.

### Mucopolysaccharidoses (MPS) and Mucopolidoses (ML)

In these groups, some exhibit significant neurologic manifestations, which depend on infiltrative deposition of MPS or oligosaccharide material into the brain, dura, or peripheral nervous system. Optic nerve infiltration can lead to blindness. One characteristic change in this group of disorders is the J-shape sella turcica produced



by the deposition of material (cystic in nature) within the cysternal spaces adjacent to the sella and eroding the bone without frank pituitary destruction [Nelson et al., 1988; Sheridan et al. 1994].

In addition, odontoid hypoplasia and C1 C2 subluxation as well as C3 C4 dislocations have been observed in some [Thomas et al., 1985]. Dural thickening at the craniocervical junction also can lead to cord compression neuropathy [Stevens et al., 1991]. Farther down the spine, the dural thickening (infiltration) may result in more peripheral neuropathy such as a neurogenic bladder as well as pain and parasthesias [Young et al., 1980]. Secondary syrinx of the cord (probably post traumatic) has been observed. Kyphosis, and even severe gibbus, is not unusual, also leading to neurologic symptomatology. Carpal tunnel syndrome can occur [Wraith et al., 1990].

**CT and MRI changes.** CT and MRI changes are quite similar in these disorders and appear to vary mostly in the severity of involvement [Lee et al., 1993; Murata et al., 1989; Taccone et al., 1993].

**CT.** Low attenuation areas occur in the brain white matter.

**MRI.** Delay in myelination may be present and high signal intensity on T2 with cystic areas scattered in many regions, which may be associated with ventriculomegaly. The anatomy of the involvement of the dura in the craniocervical region and lower spine is best delineated by this type of imaging. The same is apparent in MRI studies of the optic nerve.

This review of the neurologic findings in the skeletal dysplasias is meant to be helpful not only for the diagnosis of this large group of disorders, but also to be useful in the management of this diverse group of patients.

## ACKNOWLEDGMENTS

The author thanks M.A. Priore and S. Levin of the Skeletal Dysplasia Registry and Patti Hamm for their editorial help and patience. This study was performed in part with the assistance of NIH grant #PO1 HD 22657-08.

## REFERENCES

- Abdel-Al YK, Shabani IS, Lubani MM, al-Ghawabi MA, Ibrahim MD, al-Mohtasseb S, Duodin KI (1994): Autosomal recessive osteopetrosis in Arab children. *Ann Tropical Paediatrics* 14(1):59-64.
- Adès LC, Morris LL, Burns R, Haan EA (1994): Neurological involvement in Worth type endosteal hyperostosis: Report of a family. *Am J Med Genet* 51(1):46-50.
- Al-Mefty O, Fox JL, Al-Rodhan N, Dew JH (1988): Optic nerve decompression in osteopetrosis. *J Neurosurg* 68:80-84.
- Aleck KA, Grix A, Clericuzio C, Kaplan P, Adomian GE, Lachman R, Rimoin DL (1987): Dyssegmental dysplasias: Clinical, radiographic, and morphologic evidence of heterogeneity. *Am J Med Genet* 27(2):295-312.
- Allen HA, Haney P, Rao KC (1982): Vascular involvement in cranial hyperostosis. *AJNR* 3:193-195.
- Anderson CE, Silience DO, Lachman RS, Toomey K, Bull M, Dorst J, Rimoin DL (1982): Spondylometaphyseal dysplasia, Strudwick type. *Am J Med Genet* 13:243-256.
- Andres KH, Becker PS, Holden JK, Sharer LR, Cornford ME, Hansen LA, Hamilton R, Vinters HV (1993): Multifocal necrotizing leukoencephalopathy with pontine predilection in immunosuppressed patients. *Hum Pathol* 24:897-904.
- Applegate LJ, Applegate GR, Kemp SS (1991): MRI of multiple cranial neuropathies in a patient with Camurati-Engelmann disease: Case report. *AJNR* 12:557-559.
- Arenberg IK, Shambaugh GE Jr, Valvassori GE (1974): Otolaryngologic manifestations of frontometaphyseal dysplasia. *Arch Otolaryngol* 99:52-58.
- Aryanpur J, Hurko O, Francomano C, Wang H, Carson B (1990): Craniocervical decompression for cervicomedullary compression in pediatric patients with achondroplasia. *J Neurosurg* 73:375-382.
- Balza OR, Baza PC, Cidjl H, Pascual-Castroviejo I (1985): Neurological changes in TRP type 1. *An Esp Pediatr* 22:143-148.
- Bartynski WS, Barnes PD, Wassman JK (1989): Cranial CT of autosomal recessive osteopetrosis. *AJNR* 10:543-550.
- Benecke JE, Jr (1993): Facial nerve dysfunction in osteopetrosis. *Laryngoscope* 103:494-497.
- Canki-Klain N, Stanescu V, Stanescu R, Sinkovec J, Debevec M, Maroteaux P (1992): Lethal short limb dwarfism with dysmorphic face, omphalocele, and severe ossification defect: Piepkorn syndrome or severe boomerang dysplasia? *Ann Génét* 35:129-133.
- Charnas LR, Marini JC (1993): Communicating hydrocephalus, basilar invagination, and other neurologic features in osteogenesis imperfecta. *Neurol* 43:2603-2608.
- Chervenak FA, Isaacson G, Rosenberg JC, Kardon NB (1986): Antenatal diagnosis of frontal cephalocele in a fetus with atelosteogenesis. *J Ultrasound Med* 5:111-113.
- Chitayat D, Silver K, Azouz EM (1992): Skeletal dysplasia, intracerebral calcifications, optic atrophy, hearing impairment and mental retardation: Nosology of dysosteosclerosis. *Am J Med Genet* 43:517-523.
- Cideciyan D, Rodriguez MM, Haun RL, Abdenour GE, Bruce JH (1993): New findings in short rib syndromes. *Am J Med Genet* 46:255-259.
- Coscia MF, Bassett GS, Bowen JR, Ogilvie JW, Winter RB, Simonton SC (1989): Spinal abnormalities in campomelic dysplasia. *J Pediatr Orthop* 9:6-14.
- Coulter CL, Leech RW, Brumback RA, Schaefer GB (1991): Cerebral abnormalities in thanatophoric dysplasia. *Child's Nerv Syst* 7:21-26.
- Currarino G (1989): Primary spondylolysis of the axis vertebra (C<sub>2</sub>) in three children including one with pyknodysostosis. *Pediatr Radiol* 19:535-538.
- Demirci A, Sze G (1991): Cranial osteopetrosis: MR findings. *AJNR* 12:781-782.
- Dore DD, MacEwen GD, Boulous MI (1987): Cleidocranial dysostosis and syringomyelia. *Clin Orthop* 214:229-234.
- Dort JC, Pollak A, Fisch U (1990): The fallopian canal and facial nerve in sclerosteosis of the temporal bone. *Am J Otol* 11:320-325.
- du Plessis JJ (1993): Sclerosteosis: Neurosurgical experience with fourteen cases. *J Neurosurg* 78:388-392.
- Dvorak DM, Rusnak RA, Morcos JJ (1993): Multiple trauma in the achondroplastic dwarf. *Am J Emerg Med* 11:390-395.
- Ellis DS, Heckenlively JR, Martin CL, Lachman RS, Sakati NA, Rimoin DL (1984): Leber's congenital amaurosis associated with familial juvenile nephronophthisis and cone-shaped epiphyses of the hands (the Saldino-Mainzer syndrome). *Am J Ophthalmol* 97(2):233-239.
- Elster AD, Theros EG, Key LL, Chen MY (1992): Cranial imaging in autosomal recessive osteopetrosis: part I, II. *Radiol* 183:129-135.
- Figueiredo J, Reis A, Vaz R, Leao M, Cruz C (1989): Porencephalic cyst in pyknodysostosis. *J Med Genet* 26(12):782-784.
- Floman Y, Gomori JM, Fast A (1989): Isthmic spondylolisthesis in pyknodysostosis. *J Spinal Disorders* 2:268-271.
- Fortuna A, Ferrate L, Acqui M, Santoro A, Mastronardi L (1989): Narrowing of thoraco-lumbar spinal canal in achondroplasia. *J Neurosurg Sci* 33:185.
- Fryns JP, Van den Berghe H (1988): Facial paralysis at the age of 2 months as a first clinical sign of van Buchem disease. *Eur J Pediatr* 147:99-100.
- Gay BB, Jr., Elsas LJ, Wyly JB, Pasquali M (1994): Osteopathia striata with cranial sclerosis. *Pediatr Radiol* 24:56-60.
- Goodman P, Dominguez R (1990): Cervicothoracic myelopathy in Conradi-Hunermann disease: MRI diagnosis. *Magn Reson Imaging* 8:647-650.

- Hahn YS, Engelhard HH 3d, Naidich T, McLone DG (1989): Paraplegia resulting from thoracolumbar stenosis in a seven-month-old achondroplastic dwarf. *Pediatr Neurosci* 15:39-43.
- Hamanishi C (1993): Acrodysostosis associated with spinal canal stenosis. *Spine* 18:1922.
- Hamers A, Jongbloet P, Peeters G, Fryns JP, Geraedts J (1990): Severe mental retardation in a patient with tricho-rhino-phalangeal syndrome type I and 8q deletion. *Eur J Pediatr* 149:618-620.
- Hecht JT, Horton WA, Reid CS, Pyeritz RE, Chakraborty R (1989): Growth of the foramen magnum in achondroplasia. *Am J Med Genet* 32:528-535.
- Hecht JT, Nelson FW, Butler IJ, Horton WA, Scott CI Jr, Wassman ER, Mehlinger CM, Rimoin DL, Pauli RM (1985): Computerized tomography of the foramen magnum: Achondroplastic, values compared to normal standards. *Am J Med Genet* 20:355-360.
- Hill SC, Stein SA, Dwyer A, Altman J, Dorwart R, Doppman J (1986): Cranial CT findings in sclerosteosis. *AJNR* 7:505-511.
- Hudgins RJ, Edwards MS (1987): Craniometaphyseal dysplasia associated with hydrocephalus: Case report. *Neurosurgery* 29:617-618.
- Kahanovitz N, Rimoin DL, Silience DO (1982): The clinical spectrum of lumbar spine disease in achondroplasia. *Spine* 7:137-140.
- Karras D, Karargiris G, Vassilopoulos D, Karatzetzos C (1993): Reflex sympathetic dystrophy syndrome and osteogenesis imperfecta. *J Rheumatol* 20:162-164.
- Kirkpatrick DB, Rimoin DL, Kaitila I, Goodman SJ (1977): The cranio-tubular bone modeling disorders: A neurosurgical introduction to rare skeletal dysplasias with cranial nerve compression. *Surg Neurol* 7:221-232.
- Kitagawa Y, Hirose G, Kosoegawa H, Kanda S, Hamada S (1986): A case of van Buchem's disease with multiple cranial neuropathy. *Rinsho Shinkeigaku* 26:28-33.
- Kreck J, Starshak RJ (1987): Cervical kyphosis in diastrophic dwarfism: CT and MR findings. *Pediatr Radiol* 17:321-322.
- Lachman R, Silience D, Rimoin D, Horton W, Hall J, Scott C, Spranger J, Langer L (1981): Diastrophic dysplasia: The death of a variant. *Radiology* 140:79-86.
- LeDoux MS, Naftalis RC, Aronin PA (1991): Stabilization of the cervical spine in spondyloepiphyseal dysplasia congenita. *Neurosurgery* 28:580-583.
- Lee C, Dineen TE, Brack M, Kirsch JE, Runge VM (1993): The mucopolysaccharidoses: Characterization by cranial MR imaging. *AJNR* 14:1285-1292.
- Lenti C, Paganoni P, Sangermani R (1991): Rhizomelic chondrodysplasia punctata: 16-year follow-up of a child from birth. *Italian J Neurol Sci* 12(5):469-473.
- Lundar T, Bakke SJ, Nornes H (1990): Hydrocephalus in an achondroplastic child treated by venous decompression at the jugular foramen. *J Neurosurg* 73:138-140.
- Mador MJ, Tobin MJ (1990): Apneustic breathing: A characteristic feature of brain stem compression in achondroplasia? *Chest* 97: 877-883.
- Mah V, Nelson L, Vinters HV (1987): Focal pontine leukoencephalopathy in a patient with Shwachman-Diamond syndrome. *Can J Neurol Sci* 14:608-610.
- Martínez-Frías ML, Bermejo E, Urioste M, Egues J, Lopez Soler JA (1993): Short rib polydactyly syndrome (SRPS) with anencephaly and other central nervous system anomalies. *Am J Med Genet* 47:782-787.
- McArdle DQ, Sawaya R, Rhodadad G (1984): Brain tumor and achondroplasia: A case report and review of the literature. *Neurosurgery* 15:111-113.
- McCleary L, Rovit RL, Murali R (1987): Case report: Myelopathy secondary to congenital osteopetrosis of the cervical spine. *Neurosurgery* 20:487-489.
- Merrill KD, Schmidt TL (1989): Occipitoatlantal instability in a child with Kniest syndrome. *J Pediatr Orthop* 9:338-340.
- Murata R, Nakajima S, Tanaka A, Miyagi N, Matsuoka O, Kogame S, Inoue Y (1989): MR imaging of the brain in patients with mucopolysaccharidosis. *AJNR* 10:1165-1170.
- N'dri K, Lapra-Lepine C, Colonna C, Tavernier T, Bochu M (1994): Imaging of spinal osteochondroma of neurologic nature. *J Radiology* 75 (8-9):427-431.
- Nakayama Y, Takeno Y, Tsugu H, Tomonaga M (1994): Maffucci's syndrome associated with intracranial chordoma. *Neurosurg* 34: 907-909.
- Nelson J, Broadhead D, Mossman J (1988): Clinical findings in 12 patients with MPS IV A (Morquio's disease): Further evidence for heterogeneity. Part III: Odontoid dysplasia. *Clin Genet* 33: 111-120.
- Nöel S, Bakeriaux D, Telerman-Toppet N (1992): Unusual occurrence of cervical myelopathy in a case of Stickler's syndrome. *J Med Genet* 29:200-202.
- O'Brien MF, Bridwell KH, Lenke LG, Schoenecker PL (1994): Intracranial osteochondroma producing spinal cord compression in hereditary multiple exostoses. *J Spinal Disorders* 7(3):236-241.
- Paludetti G, Galli A, Lombardi R, Almadori G, Pujolas FX, Maurizi M. (1994): Otolaryngological aspects of Camurati-Engelmann disease (progressive diaphyseal dysplasia): Review of literature and report of one case. *Acta Otorrinolaringologica Espanola*, 45(3):207-213.
- Popovic-Rolovic M, Calic-Perisic N, Bunjevacki G, Negovanovic D (1976): Juvenile nephronophthisis associated with retinal pigmentary dystrophy, cerebellar ataxia, and skeletal abnormalities. *Arch Disease in Childhood* 51(10):801-803.
- Pauli RM, Gilbert EF (1986): Upper cervical cord compression as cause of death in osteogenesis imperfecta type 2. *J Pediatr* 108: 579-581.
- Pauli RM, Horton VK, Glinski LP, Reiser CA (1995): Prospective assessment of risks for cervicomedullary-junction compression in infants with achondroplasia. *Am J Human Genet* 56(3):732-744.
- Pauli RM, Scott CI, Wassmas ER Jr, Gilbert EF, Leavitt LA, Ver Hoeve J, Hall JG, Partington MW, Jones KL, Sommer A, et al (1984): Apnea and sudden unexpected death in infants with achondroplasia. *J Pediatr* 104:342.
- Phadke RV, Phadke SR, Chakravarti AL (1990): Spinal dysraphism in achondroplasia. *Pediatr Neurosurg* 9:32-34.
- Prudio J, Stollenburg-Didinger G, Jimenez E, Cervos-Navarro J (1993): Central nervous system alterations in a case of short rib polydactyly syndrome Majewski type. *Devel Med Child Neurology* 35:158.
- Rakic M, Elhosseiny A, Ramadan F, Iyer R, Howard RG, Gross L (1986): Adult-type osteopetrosis presenting as carpal tunnel syndrome. *Arthritis Rheum* 29:926-928.
- Ramer JC, Eggl K, Rogan PK, Laadda RL (1993): Identical twins with Weissbacher-Zweymüller syndrome and neural tube defects. *Am J Med Genet* 45:614-618.
- Reardon W, Hall CM, Shaw DG, Kendall B, Hayward R, Winter RM (1994): New autosomal dominant form of spondyloepiphyseal dysplasia presenting with atlanto-axial instability. *Am J Med Genet* 52(4):432-437.
- Reid CS, Pyeritz RE, Kopits SE, Maria BL, Wang H, McPherson RW, Hurko O, Phillips JA 3d, Rosenbaum AE (1987): Cervicomedullary compression in young patients with achondroplasia. *J Pediatr* 110:522-530.
- Richards BS (1991): Atlanto-axial instability in diastrophic dysplasia. *JBJS* 73A:614-616.
- Robinson D, Tieder M, Halperin M, Burshtein D, Nevo Z (1994): Maffucci's syndrome—the result of neural abnormalities? *Cancer* 74:949-957.
- Rosenberg E, Lohr H (1986): A new hereditary bone dysplasia with characteristic bowing and thickening of the distal ulna. *Eur J Pediatr* 145:40-45.
- Rush GA, Burke SW (1984): Hangman's fracture in a patient with osteogenesis imperfecta. *JBJS* 66A:778-779.
- Saper JR, Holt JF (1974): Cranial metaphyseal dysplasia: A cause of recurrent bilateral facial palsy. *Arch Neurol* 31:204-207.
- Sheridan M, Johnston I (1994): Hydrocephalus and pseudotumour cerebri in the mucopolysaccharidoses. *Child's Nerv Syst* 10: 148-150.
- Shohat M, Lachman R, Rimoin DL (1989): Odontoid hypoplasia, cervical spine subluxation and hydrocephalus in metatropic dysplasia. *J Pediatr* 114:239-243.
- Silience DO (1994): Craniocervical abnormalities in osteogenesis imperfecta. *Pediatr Radiol* 24:427-430.
- Singh M, Ray D, Paul VK, Kumar A (1988): Hydrocephalus in asphyxiating thoracic dystrophy. *Am J Med Genet* 29:391-395.

- Spranger JW, Bierbaum B, Herrmann J (1976): Heterogeneity of Dyggre-Melchior-Clausendwarfism. *Hum Genet* 33:279-287.
- Steinbok P, Hall J, Flodmark O (1989): Hydrocephalus in achondroplasia: The possible role of intracranial venous hypertension. *J Neurosurg* 71:42-48.
- Steinsapir KD, Vinters HV (1985): Central pontine myelinolysis in a child with Shwachman-Diamond syndrome. *Hum Pathol* 16:741-743.
- Stevens JM, Kendall BE, Crockard HA, Ransford A (1991): The odontoid process in Morquio-Brailsford's disease: The effects of occipitocervical fusion. *J Bone Joint Surg (Br)* 73B:851-858.
- Taccone A, Tortori Donati P, Marzoli A, Dell'Acqua A, Gatti R, Leone D (1993): Mucopolysaccharidosis: Thickening of dura mater at the craniocervical junction and other CT/MRI findings. *Pediatr Radiol* 23:349-452.
- Taybi H, Lachman RS (1996): "Radiology of Syndromes, Metabolic Disorders and Skeletal Dysplasias," 4th ed. St. Louis: Mosby-Yearbook.
- Thomas SL, Childress MH, Quinton B (1985): Hypoplasia of the odontoid with atlanto-axial subluxation in Hurler's syndrome. *Pediatr Radiol* 15:353.
- Trabelsi M, Hammou-Jeddi A, Kammoun A, Bennaceur B, Gharbi HA (1990): Asphyxiating thoracic dysplasia with hepatic ductal hypoplasia, agenesis of the corpus callosum and Dandy Walker malformation. *Pédiatrie* 45:35-38.
- Tsipouras P, Barabas G, Matthews WS (1986): Neurologic correlates of osteogenesis imperfecta. *Arch Neurol* 43:150-152.
- Waters KA, Everett F, Sillence D, Fagan E, Sullivan CE (1993): Breathing abnormalities in sleep in achondroplasia. *Arch Dis Childh* 69:191-196.
- Weisz GM (1982): Lumbar spinal canal stenosis in osteopoikilosis. *Clin Orthop* 166:89-92.
- Wieting JM, Krach LE (1994): Spinal cord injury rehabilitation in a pediatric achondroplastic patient: Case report. *Arch Physical Medicine and Rehabilitation* 75(1):106-108.
- Wilms G, Casaer P, Alliet P, Demaerel P, Smet M, Baert AL (1990): Cerebrovascular occlusive complications in osteopetrosis major. *Neuroradiol* 32:511-513.
- Wraith JE, Alani SM (1990): Carpal tunnel syndrome in the mucopolysaccharidoses. *Arch Dis Child* 65:962-963.
- Yagi Y, Yamazaki M, Satone T (1987): Anesthesia for patients with spondyloepiphyseal dysplasia congenita. *Masui* 36:793-796.
- Yamada H, Nakamura S, Tajima M, Kageyama N (1981): Neurological manifestations of pediatric achondroplasia. *J Neurosurg* 54:49-57.
- Yamamoto T, Tohyama J, Koeda T, Maegaki Y, Takahashi Y (1995): Multiple epiphyseal dysplasia with small head, congenital nystagmus, hypoplasia of corpus callosum, and leukonychia totalis: A variant of Lowry-Wood syndrome? *Am J Med Genet* 56(1):6-9.
- Yamashita S, Izawa T, Misugi N, Sasaki Y (1994): A case of motor and sensory neuropathy associated with metatropic dysplasia. *No To Hattatsu Brain and Development* 26(3):258-262.
- Young R, Kleinman G, Ojemann RG, Kolodny E, Davis K, Halperin J, Zalneraitis E, DeLong GR (1980): Compressive myelopathy in Maroteaux-Lamy syndrome: Clinical and pathological findings. *Ann Neurol* 8:336-340.
- Zonana J, Rimoin DL, Lachman RS, Cohen AH (1977): A unique chondrodysplasia secondary to a defect in chondroosseous transformation. *Birth Defects Original Article Series* 13(3D):155-163.